



Genomic Sequence of a Ranavirus Isolated from Short-Finned Eel (*Anguilla australis*)

Subramaniam, Kuttichantran; Toffan, Anna; Cappellosza, Elisabetta; Steckler, Natalie K; Olesen, Niels Jørgen; Ariel, Ellen; Waltzek, Thomas B

Published in:
Genome Announcements

Link to article, DOI:
[10.1128/genomeA.00843-16](https://doi.org/10.1128/genomeA.00843-16)

Publication date:
2016

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Subramaniam, K., Toffan, A., Cappellosza, E., Steckler, N. K., Olesen, N. J., Ariel, E., & Waltzek, T. B. (2016). Genomic Sequence of a Ranavirus Isolated from Short-Finned Eel (*Anguilla australis*). *Genome Announcements*, 4(4), [e00843-16]. <https://doi.org/10.1128/genomeA.00843-16>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Genomic Sequence of a Ranavirus Isolated from Short-Finned Eel (*Anguilla australis*)

 Kuttichantran Subramaniam,^a Anna Toffan,^b Elisabetta Cappelozza,^b Natalie K. Steckler,^a Niels J. Olesen,^c Ellen Ariel,^d Thomas B. Waltzek^a

Department of Infectious Diseases and Pathology, College of Veterinary Medicine, University of Florida, Gainesville, Florida, USA^a; National Reference Laboratory for Fish Diseases, Istituto Zooprofilattico Sperimentale delle Venezie, Legnaro, Italy^b; National Veterinary Institute, Technical University of Denmark, Frederiksberg C, Denmark^c; College of Public Health, Medical and Veterinary Sciences, James Cook University, Townsville, Queensland, Australia^d

The short-finned eel ranavirus (SERV) was isolated from short-finned eel imported to Italy from New Zealand. Phylogenomic analyses revealed that SERV is a unique member of the genus *Ranavirus*, family *Iridoviridae*, branching at the base of the tree near other fish ranaviruses.

Received 21 June 2016 Accepted 23 June 2016 Published 18 August 2016

Citation Subramaniam K, Toffan A, Cappelozza E, Steckler NK, Olesen NJ, Ariel E, Waltzek TB. 2016. Genomic sequence of a ranavirus isolated from short-finned eel (*Anguilla australis*). *Genome Announc* 4(4):e00843-16. doi:10.1128/genomeA.00843-16.

Copyright © 2016 Subramaniam et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Thomas B. Waltzek, tbwaltzek@ufl.edu.

Short-finned eel ranavirus (SERV) was isolated in 1999 from short-finned eel (*Anguilla australis*) imported to Italy from New Zealand (1). Ten apparently healthy eels weighing between 200 and 300 g were processed for diagnostic virology as part of a routine screening of imported live fish. Internal organ homogenates seeded onto epithelioma papulosum cyprini (EPC) cells resulted in CPE characterized by focal plaques. Negative stain electron microscopy performed on infected culture supernatant revealed iridovirus-like particles. Infected cell cultures reacted positively in a direct immunofluorescence assay using a rabbit polyclonal serum against the fish ranavirus, *European catfish virus* (ECV, [2]), suggesting that the replicating agent was a member of the genus *Ranavirus*.

A sixth passage of the SERV isolate was amplified in EPC cells maintained in modified Eagle's medium with 10% fetal bovine serum at 23°C. Inoculation of EPC cells at high multiplicity of infection provided material from an eighth passage harvested after 96 h. Cell culture supernatant was clarified at 3,000 × g for 20 min, and total nucleic acids were purified using a DNeasy blood and tissue kit (Qiagen). A DNA library was prepared using a Nextera XT DNA kit (Illumina) and sequenced using a V3 chemistry 600-cycle kit on a MiSeq platform (Illumina). *De novo* assembly of 10,752,536 paired-end reads in SPAdes (3) produced a contiguous consensus sequence of 126,965 bp with a G+C content of 55.64%. The quality of the genome assembly was assessed by mapping the reads back to the consensus sequence in Bowtie 2 (4) and visually inspecting the alignment in Tablet (5). A total of 6,417,927 reads (59.69%) aligned at an average coverage of 10,720 reads per nucleotide.

The genome of SERV was annotated using GATU (6) with *Frog virus 3* (GenBank accession no. NC_005946) as the reference. Additional putative open reading frames (ORFs) were identified using GenemarkS (7). A total of 111 putative ORFs were predicted in SERV compared to other related fish ranaviruses, including 100 in *Epizootic hematopoietic necrosis virus* (EHNV, GenBank

accession no. NC_028461), 135 in ECV (GenBank accession no. KT989884), and 136 in European sheatfish virus (ESV, GenBank accession no. JQ724856). An analysis of locally collinear blocks in Mauve (8) revealed that the genomes of SERV, EHNV, ECV, and ESV are collinear. Phylogenetic analyses based on the concatenated nucleotide sequences of the 26 *Iridoviridae* core genes (9) revealed that SERV forms a distinct branch at the base of the ranavirus tree near other fish ranaviruses (e.g., EHNV, ESV, and ECV) with only the highly divergent Santee-Cooper ranavirus and grouper iridoviruses splitting off earlier.

Although some fish ranaviruses can cause considerable mortality in aquaculture (e.g., EHNV, ECV, and grouper iridoviruses), others like SERV are rarely detected and thus their impacts are unknown (10). Bath challenges with SERV resulted in significant mortality in northern pike *Esox lucius* fry (11) versus no appreciable disease in juvenile black bullheads *Ameiurus melas* (12). Thus, SERV appears to have minimal impacts on the health of some hosts, including short-finned eel, while causing disease in others.

Accession number(s). The complete genome sequence of SERV has been deposited in GenBank under accession number [KX353311](https://www.ncbi.nlm.nih.gov/nuclot/KX353311).

ACKNOWLEDGMENTS

We thank Giuseppe Bovo for his dedicated and influential work on fish viruses, including SERV. We also thank Federica Gobbo and Patrick Thompson for their technical assistance.

FUNDING INFORMATION

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

REFERENCES

1. Bovo G, Giacometti G, Montesi F, Cappelozza E, Ormelli S. 1999. Isolation of an irido-like viral agent from New Zealand eel. Poster 153. 9th International Conference of European Association of Fish Pathologists, Rhodes, Greece.

2. Bovo G, Comuzzi M, De Mas S, Ceschia G, Giorgetti G, Giacometti P, Cappellozza E. 1993. Isolation of an irido-like viral agent from breeding catfish (*Ictalurus melas*). Bollettino Società Italiana di Patologia Ittica 11: 3–10.
3. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. J Comput Biol 19:455–477. <http://dx.doi.org/10.1089/cmb.2012.0021>.
4. Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nat Methods 9:357–359. <http://dx.doi.org/10.1038/nmeth.1923>.
5. Milne I, Bayer M, Cardle L, Shaw P, Stephen G, Wright F, Marshall D. 2010. Tablet—next generation sequence assembly visualization. Bioinformatics 26:401–402. <http://dx.doi.org/10.1093/bioinformatics/btp666>.
6. Tcherepanov V, Ehlers A, Upton C. 2006. Genome annotation transfer utility (GATU): rapid annotation of viral genomes using a closely related reference genome. BMC Genomics 7:150. <http://dx.doi.org/10.1186/1471-2164-7-150>.
7. Besemer J, Lomsadze A, Borodovsky M. 2001. GeneMarkS: a self-training method for prediction of gene starts in microbial genomes: implications for finding sequence motifs in regulatory regions. Nucleic Acids Res 29:2607–2618. <http://dx.doi.org/10.1093/nar/29.12.2607>.
8. Darling AC, Mau B, Blattner FR, Perna NT. 2004. Mauve: multiple alignment of conserved genomic sequence with rearrangements. Genome Res 14:1394–1403. <http://dx.doi.org/10.1101/gr.2289704>.
9. Eaton HE, Metcalf J, Penny E, Tcherepanov V, Upton C, Brunetti CR. 2007. Comparative genomic analysis of the family Iridoviridae: re-annotating and defining the core set of iridovirus genes. Virol J 4:11. <http://dx.doi.org/10.1186/1743-422X-4-11>.
10. Duffus ALJ, Waltzek TB, Stöhr AC, Allender MC, Gotesman M, Whittington RJ, Hick P, Hines MK, Marschang RE. 2015. Distribution and host range of ranaviruses, p. 9–57. In Gray MJ, Chinchar VG (ed), Ranaviruses: lethal pathogens of ectothermic vertebrates. Springer, New York. http://dx.doi.org/10.1007/978-3-319-13755-1_2.
11. Bang-Jensen BB, Ersbøll AK, Ariel E. 2009. Susceptibility of pike *Esox lucius* to a panel of ranavirus isolates. Dis Aquat Organ 83:169–179. <http://dx.doi.org/10.3354/dao02021>.
12. Gobbo F, Cappellozza E, Pastore MR, Bovo G. 2010. Susceptibility of black bullhead *Ameiurus melas* to a panel of ranavirus isolates. Dis Aquat Organ 90:167–174. <http://dx.doi.org/10.3354/dao02218>.